

Accelerator R&D in Europe

Subject → Advanced Accelerator R&D



R. Assmann

Leading Scientist DESY

Coordinator Preparation Team

Helmholtz Distributed ARD Test Facility

Coordinator European Network for Novel Accelerators
(EuroNNAc)


Deputy-Coordinator European Coordination for
Accelerator R&D (EuCARD2)

Accelerator R&D HEPAP Subpanel
SLAC, 29.8.2014




- > **Accelerators at the Energy Frontier**
- > Accelerator Builder's Challenge
- > EuPRAXIA – Proposal for a European Plasma Accelerator for Users
- > Helmholtz ARD Program
- > Conclusions

Livingston and Accelerators at the Energy Frontier




Why this saturation?

1. Conventional accelerators have reached **hard physical limits**.
2. Energy is extended by scaling or modestly extending proven accelerator technology. As a consequence, **length and project cost also scale with energy**.
3. Ever bigger projects require more and more time to bring.




Livingston and Accelerators at the Energy Frontier



Livingston and Accelerators at the Energy Frontier

ILC project for the 2020's/2030's. FCC project for the 2030's/2040's.


1. Timelines reflect the size, complexity and cost of projects (scaling effects).
2. These projects will be very mature and their performances cannot easily be matched by novel accelerator technologies.
3. However, it is possible that the required funding for ILC or FCC cannot be obtained.
4. In this case it will be crucial to have a plan B HEP accelerator project in the 2030's.
 - a. This project will likely be less performant than ILC.
 - b. This project should also be much smaller and much lower cost.




ar



Livingston and Accelerators at the Energy Frontier




Livingston and Accelerators at the Energy Frontier




Advent of plasma acc.:


1. Metallic cavity walls replaced with plasma walls → overcoming hard physical limits of metallic RF structures.
2. Acceleration lengths (same energy) are **factor 100 – 1000 shorter**. Multi-GeV e-beams proven.
3. Still short-comings but **no fundamental limit**.
4. Financial effort in HEP at or below 1% of total. **More resources will lead to faster progress**.




Livingston and Accelerators at the Energy Frontier




Livingston and Accelerators at the Energy Frontier



Livingston and Accelerators at the Energy Frontier




Livingston and Accelerators at the Energy Frontier



- > Accelerators at the Energy Frontier
- > **Accelerator Builder's Challenge**
- > EuPRAXIA – Proposal for a European Plasma Accelerator for Users
- > Helmholtz ARD Program
- > Conclusions


Reminder: Plasma-Acceleration (Internal Injection)

Laser Pulse (200 TW, ~ 30 fs, $E_{\text{transv}} \sim \text{TV/m}$)




Plasma electrons
(plasma cell, $\sim 10^{19} \text{ cm}^{-3}$)


Reminder: Plasma-Acceleration (Internal Injection)



Reminder: Plasma-Acceleration (Internal Injection)




Reminder: Plasma-Acceleration (Internal Injection)





This accelerator fits into a human hair!

Reminder: Plasma-Acceleration (Internal Injection)



- This proved highly successful with electron bunches of **up to 4.25 GeV produced over a few cm.**
- Small dimensions involved \rightarrow few **micron tolerances!**
- Highly compact but also **highly complex** accelerator: generation, bunching, focusing, acceleration, (wiggling) all in one small volume.
- Energy spread and stability at the few % level.

External Injection...



**Less complex.
Way to acceleration
stages.**


Accelerator Builder's Challenge *(simplified to typical values)*

- > Match into and out of plasma with **beam size around $1\ \mu\text{m}$** (about 1 mm beta function).
- > Control **offsets** between the wakefield driver (laser or beam) and the accelerated electron bunch at **$1\ \mu\text{m}$ level**.
- > Use **short bunches (few fs)** to minimize energy spread.
- > Achieve **synchronization stability of few fs** from injected electron bunch to wakefield (energy stability and spread).
- > Control the **charge and beam loading** to compensate energy spread (idea Simon van der Meer).
- > Develop and demonstrate **user readiness of a 5 GeV plasma accelerated beam**.



Accelerator Builder's Challenge – Feasible?


- Difficult but we believe solutions can be found. Will not come for free...



50 nm with a
1.3 GeV
electron beam
(from K. Kubo et al.
Proc. IPAC 2014)

Accelerator Builder's Challenge – Feasible?

- Difficult but we believe solutions can be found. Will not come for free...



Femtosecond Precision in Laser-to-RF Phase Detection

(from H. Schlarb, T. Lamb, E. Janas et al. Report on DESY Highlights 2013).

- Again: **No fundamental limit here, but strong technical challenges!**

- > Accelerators at the Energy Frontier
- > Accelerator Builder's Challenge
- > **EuPRAXIA – Proposal for a European Plasma Accelerator for Users**
- > Helmholtz ARD Program
- > Conclusions

- Quite a number of national investments in the EU member states at the 10 – 50 M€ level.
 - Europe is very competitive: excellent lasers, large expertise, big teams with many young students/researchers, new ideas from Europe have leading influence.
 - E.g. invention of “bubble regime” in Europe:

Pukhov, Alexancer, and Jürgen Meyer-ter-Vehn.
"Laser wake field acceleration: the highly non-linear broken-wave regime." Applied Physics B 74.4-5 (2002): 355-361.

736 citations

SCAPA



Lund
Laser Centre



STFC
ASTeC

LAOLA

Laboratory for Laser- and beam-driven plasma Acceleration

ILPP

JuSPARC

ELBE



HELMHOLTZ
ZENTRUM DRESDEN
ROSSENDORF



STFC

Central Laser Facility

Cilex

Centre Interdisciplinaire Lumière Extrême



PHELIX

eli



beamlines

ALA



Laboratoire d'optique appliquée

UMR 2039 - Palaiseau - France

AIVAKE



Consiglio
Nazionale delle
Ricerche

SPARC



INFN
Sezione di Roma



- > The big French project.

5 PW laser and LWFA area




Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 17




- > Well established lab, since 1992.


High stability LWFA



Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 22





X-rays



- > Electron beam: 150 MeV, multi-bunch, bunch length below 300 fs, 200 pC, 1 μ m norm. emittance
- > FLAME laser: Ti:Sa, chirped pulse amplification (CPA), 200 TW, 25 fs long, 10 Hz repetition rate
- > SPARC: Incoherent plasma wakefield experiment

Comb beam \rightarrow high efficiency

- > Comb project is unique: resonant beam driven plasma wakefields.



Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 21




- > Salle Jaune Laser: 70 TW, repetition rate 10 Hz, pulse duration of 30 fs

- > Goals:

- Exploration of new laser plasma accelerator physics for the production of electron beams

LWFA for FEL



Victor Malka
Researcher at CNRS and Lecturer
An excellence grant for LOA.



Victor Malka is a CNRS researcher and lecturer in the physics department at X, works at ENSTA. In a team that he set up in 2001 to study laser-plasma particle acceleration. In July 2008, he was awarded a grant by the European Research Council of 2.2 million euros. The grant was awarded in two categories (junior and senior) it was in the second category that he was rewarded for his many scientific works and for his ability to create new fields of research.




Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 18



- > Builds on expertise at MPQ and LMU.
- > Successful in laser-driven generation of electron beams. See example with MPQ lasers.
- > Ongoing: 63 M€ investment. Financed by State Bavaria and CAI 201

LWFA for science (FEL, ...)

from gas cell: 600 MeV, 200 pC
X experiments into water window



Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 26



- > Laser: 10-15 fs duration, up to 10 PW. End stage: a few kJ in 15 fs (~200 PW) with low repetition rate (minute based).

- > Might be

- > New test display tools for testing,

- > Laser-driven and pre-unprec 100 GeV for the quality cancer therapy.

10 – 200 PW laser, also for LWFA (finally 100 GeV?)



beamlines



Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 23

- > COXINEL: COherent Xray source INferred from Electrons accelerated by Laser.

- > Leader: Marie-Emmanuelle Couprie, SOLEIL.

- > Goals:

- COXINEL is a new FEL R&D for LWFA
- FEL R&D for LWFA
- FEL R&D for LWFA


FEL R&D for LWFA

acceleration, it
mplification

intense light



Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 19



ICAN for high efficiency

Figure 1 | Protons (Z), and split stages produced compressed -10 kHz (Z).

The future is fibre accelerators

Gerard Mourou, Bill Brocklesby, Toshiki Tajima and Jens Limpert
Could massive arrays of thousands of fibre lasers be the driving force behind next-generation particle accelerators? The International Coherent Amplification Network project believes so and is currently performing a feasibility study.

Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 20



Coherent Amplification Network

- > International collaboration with approved experiment at CERN beam.
- > Driver: 450 GeV proton bunch, $1e11$, 3.5 μ m emittance, bunch length \gg plasma wavelength
- > Microbunching

Proton-driven PWFA

- 1) Protons at focusing regions survive.
 - 2) Protons at defocusing regions get lost.
- Surviving microbunches induce wakefields.
 - Accelerate injected electrons from several 10 MeV to GeV.



Ralph Admanen | UK Plasma Acc. Workshop | 31.1.2014 | Page 24





LWFA FEL

repetition rate
towards FEL.



e- driven PWFA

Beam-driven plasma wakefields. Beam-driven plasma wakefields with shaped beams and innovative injection methods.

forward




PWFA modulation

31.1.2014 | Page 32



The LIGHT test-stand at GSI: coupling of laser-accelerated ions into conventional accelerators

Principle: manipulation of laser-accelerated ions



Ion plasma acc. and transport

Cockcroft Institute: Universities of Lancaster, Liverpool and Manchester, the Science and Technology Facilities Council (STFC) and the Natural Environment Research Council (NERC).

Ultra-short CLARA project
Training.

FEL, industrial applications, PWFA



J. Clarke



C. Weis



S. Smith



Ralph Admann | Accelerator R&D Hub



ARD Spitzenforschung

LWFA low density, external inj. atto-s radiation sources




Physikalisches Institut, Universität Erlangen-Nürnberg

Ralph Admann | SINBAD | 24.01.2014 | Page 16



Helmholtz-Institute Jena



plasma wakefield imaging

ELBE center for high power radiation sources

Dual beam
Diode pumped
Synchron
Dedicated
Beam driven

Two 1 PW laser, ion/p plasma acc., radiation therapy R&D



Penelope diode pumped PW



LWFA, polarized particles



Material research

Markus Büscher

STFC – CLF and John Adams Institute in UK

STFC Central Laser Facility used also for LWFA.

John Adams Institute for Accelerator Science, Royal Holloway College, London

Plasma energy

Training of accelerator specialists, also in advanced methods.



S. Hooker



Z. Najmudin



A. Seng



R. Wulfsberg

Ralph Admann | Accelerator R&D HEPAT Subproject | 29.8.2014 | Page 66



Strathclyde in UK/Scotland



SCAPA = Scottish Centre for the Application of Plasma-based Accelerators

LWFA for generation of particle beams (electrons, protons, neutrons, etc.)

LWFA for radiation sources

Dedicated to the Production and Application of Ultra-short Electron Bunches and Radiation Pulses.



D. Jaroszynski




Z.M. Sheng

Ralph Admann | Accelerator R&D HEPAT Subproject | 29.8.2014 | Page 65



EU Funded Network on Novel Accelerators


1st European
Advanced
Accelerator
Workshop
EAAC2013




The EuPRAXIA Project Proposal

Proposal for a H2020 Design Study “European Plasma Research Accelerator with eXcellence In Applications” (EuPRAXIA^{*})

Collaborative Project



personification of well-being. In medicine the term Eupraxia
ability to perform coordinated movements.



EuPRAXIA – Connected Labs and Institutes

List of participants:

Participant no.	Participant organisation name	Short name	Country
1 (Coordinator)	Stiftung Deutsches Elektronen Synchrotron	DESY	Germany
2	Istituto Nazionale di Fisica Nucleare	INFN	Italy
3	Consiglio Nazionale delle Ricerche	CNR	Italy
4	Centre National de la Recherche Scientifique	CNRS	France
5	University of Strathclyde	USTRAH	UK
6	Instituto Superior Técnico	IST	Portugal
7	Science & Technology Facilities Council	STFC	UK
8	Synchrotron SOLEIL – French National Synchrotron	SOLEIL	France
9	University of Manchester	UMAN	UK
10	University of Liverpool	ULIV	UK
11	Agenzia nazionale per le nuove tecnologie, l'energia e lo sviluppo economico sostenibile	ENEA	Italy
12	Commissariat à l'Énergie Atomique et aux énergies alternatives	CEA	France
13	Sapienza Università di Roma	UROM	Italy
14	Universität Hansestadt Hamburg	UHH	Germany
15	University of Oxford	UOXF	UK
16	Imperial College London	ICL	UK

16 beneficiaries from 5 EU member states
plus 18 associated partners

Associated partner organisation name	Short name	Country
Jiaotong-Universität Shanghai	JUS	China
Tsinghua University Beijing	TUB	China
Extreme Light Infrastructures - Beams	ELI-B	Czech Repu
Lille University	PHLAM	France
Helmholtz Institute Jena	HIJ	Germany
Helmholtz-Zentrum Dresden-Rossendorf	HZDR	Germany
Ludwig-Maximilians-Universität München	LMU	Germany
Wigner Research Center of the Hungarian Academy of Science	WIGNER	Hungary
European Organization for Nuclear Research	CERN	IEIO ¹
High Energy Accelerator Research Organization	KEK	Japan
Kansai Photon Science Institute, Japan Atomic Energy Agency	KPSI-JAEA	Japan
Osaka University	OU	Japan
RIKEN SPring-8 Center	RSC	Japan
Lund University	LU	Sweden
Center for Accelerator Science and Education at Stony Brook U & BNL	CASE	USA
Lawrence Berkeley National Laboratory	LBNL	USA
SLAC National Accelerator Laboratory	SLAC	USA
University of California, Los Angeles	UCLA	USA



EuPRAXIA – Support

European Steering Group for Accelerator R&D, 2014

EuPRAXIA

Producing high acceleration gradients is a critical issue for particle accelerators, as highlighted by the European Strategy for Particle Physics. The feasibility of large acceleration gradient (up to 100 GV/m) within exited plasma channels is demonstrated since several decades. More recently, it has been shown that appropriate beam properties (low emittance and beam energy dispersion as well as acceptable bunch charge) can be obtained. Europe is at the forefront of this research. Based on these achievements, the realization of accelerators with appropriate beam characteristics for user-communities is now credible and highly desirable.

The vast majority of accelerators operating in the world are relatively low energy (<~10 GeV) facilities including light sources, medical and industrial applications. However these infrastructures remain complex to develop and operate and necessitate relatively large footprints.

The EuPRAXIA proposal aims at establishing the design of a 1 to 5 GeV electron accelerator with pilot applications for the Free Electron Laser user community as well as the community developing state-of-the-art particle detectors.

The proposal is technically strong and federates the major European competences and institutes required to accomplish the needed tasks.

“... the realization of accelerators with appropriate beam characteristics for user-communities is now credible and highly desirable.”

competition from CERN FCC and
ESS neutrino upgrade DS proposals Ralph A

THALES

THALES OPTRONIQUE
2 avenue Gay-Lussac – CS 90502
78995 Elancourt Cedex
France
Tél.: +33 (0)1 30 96 70 00
Fax: +33 (0)1 30 96 75 50

To whom it may concern



VF de certifié 351057

Date: 05/08/2014

Réf. /TOSA/DSL/05/08/2014

Objet : / EuPRAXIA
Subject:

Madame, Sir,

We have been informed by Mr François MATHIEU, Project Manager of APOLLON program at CNRS, about the EuPRAXIA proposal of Design Study for a Large Research Infrastructure based on laser-plasma acceleration.

Thales Optronique is recognized as leader of Intense Laser System for particles acceleration. In 2012, Thales Optronique has successfully installed at Berkeley the first commercial PetaWatt laser System for the BELLA project of US Department of Energy which is using the same physics concepts. In 2013, Thales Optronique installed a second PetaWatt laser system in Bucharest (Romania).

Thales is also involved in other large projects aiming to introduce the laser technology in the world of accelerators as it has been the case for a system now running at DESY in Germany and for another system in construction to be installed in Japan on the SACLA infrastructure involving a large free-electrons laser.

All these new projects exploit the specific advantages provided by the technology of laser-plasma acceleration and in particular the compactness inherent to the concept.


Therefore it is expected by the whole community of accelerators that these new machines will have also much better capabilities to develop societal applications like in medicine for cancer therapy and in industry for high resolution non-destructive inspection with X-rays or Gamma rays.

Therefore Thales Optronique is highly supportive of EuPRAXIA proposal for allowing European Union to be at the peak of this science and of related industry.


Sincerely yours,

Denis LEVAILLANT
Director of Laser Business Unit

Ansatz: Required Intermediate Step to Plasma LC




Schematic Layout EuPRAXIA Research Infrastructure



Present Laser
Plasma Acce-
lerators


Up to 4.25 GeV
electron beams

Schematic Layout EuPRAXIA Research Infrastructure




Schematic Layout EuPRAXIA Research Infrastructure

PLASMA ACCELERATOR




Schematic Layout EuPRAXIA Research Infrastructure

PLASMA ACCELERATOR




HEP USER AREA




EuPRAXIA
Research
Infrastructure

Schematic Layout EuPRAXIA Research Infrastructure

PLASMA ACCELERATOR



HEP USER AREA



EuPRAXIA
Research
Infrastructure

FEL / RADIATION SOURCE USER AREA

EuPRAXIA Research Infrastructure Goal Parameters

Beam Parameter	Unit	Value
Particle type	-	Electrons
Energy	GeV	1 – 5
Charge per bunch	pC	1 – 50
Repetition rate	Hz	10
Bunch duration	fs	0.01 - 10
Peak current	kA	1 – 100
Energy spread	%	0.1 – 5
Norm. emittance	mm	0.01 – 1




Envisaged Implementation


- > Goal is to design one operational facility at one location.
- > Resources will be distributed to all partners:
 - Model of big particle physics detector: Many institutes team up to build one detector at one place, each contributing a part.
- > Site study with the goal to propose the best site:
 - Existing infrastructure, host lab support, scientific user community, support from funding agency, ...
- > Facility will be devoted to provide for pilot users:
 - Ultra-compact X-ray FEL
 - Ultra-compact GeV electron source for HEP detector development



Timelines



Timelines



- > Accelerators at the Energy Frontier
- > Accelerator Builder's Challenge
- > EuPRAXIA – Proposal for a European Plasma Accelerator for Users
- > **Helmholtz ARD Program**
- > Conclusions

Research field Matter in the Helmholtz Association

 **Helmholtz-Zentrum Geesthacht**
Centre for Materials and Coastal Research

 **JÜLICH**
FORSCHUNGSZENTRUM


+ Helmholtz Institutes
Jena and Mainz


Karlsruher Institut für Technologie



Hamburg and Zeuthen

 **HZB** Helmholtz
Zentrum Berlin

 **HZDR**

 **HELMHOLTZ**
ZENTRUM DRESDEN
ROSSENDORF

since 2011

**(Total Helmholtz: 18 centers,
6 research fields, ~3B€ yearly
budget)**



Research Field Matter: new programme structure

Matter and the Universe

Fundamental Particles and Forces

Cosmic Matter in the Laboratory

Matter and Radiation from the Universe

LK II
„performance category II“
= user operation of large scale facilities

From Matter to Materials and Life

In-House Research on the Structure, Dynamics and Function of Matter at Large Scale Facilities

Facility Topic:
Research on Matter with Brilliant Light Sources

Facility Topic:
Neutrons for Research on Condensed Matter

Facility Topic:
Physics and Materials Science with Ion Beams

Facility Topic:
Research at Highest Electromagnetic Fields

Matter and Technologies

Accelerator Research and Development

Detector Technologies and Systems

Evaluation for the programme oriented funding 2015 – 19 in the research field Matter was recently completed



Accelerator R&D Programme



*Implementation phase in
Helmholtz since 2011*

ARD team at the rehearsal for the
PoF3-evaluation (HZDR, 25Feb 2014)



- > Evaluation of Research Programme MT with topics ARD and DTS
26 - 28 March, 2014
- > Whole program and ARD in particular received excellent comments in oral close-out from evaluation committee
- > Strategy with good balance between activities aiming at shorter term, direct impact on research infrastructures and more generic, visionary ones was especially praised

Participating
Centres:

Deutsches Elektronen-Synchrotron DESY
Forschungszentrum Jülich
GSI Helmholtzzentrum für Schwerionenforschung
Helmholtz-Zentrum Berlin für Materialien und Energie
Helmholtz-Zentrum Dresden-Rossendorf
Karlsruher Institut für Technologie

Programme
Speaker:


Ties Behnke (DESY)




Past and future resources in accelerator R&D

(yearly increase from 2016 still to be decided)


FTEs in Acc R&D Activities at Helmholtz centers




ARD FTEs at Helmholtz centers



3rd party funding at partner Universities



ARD full cost at Helmholtz centers



ARD structure and coordination

Topic 1: Accelerator Research and Development ARD

Speaker: R. Brinkmann/DESY, co-speaker: A. Jankowiak/HZB

ST1: SRF science and technology

Coordination:
J. Knobloch/HZB,
P. Michel/HZDR

ST2: concepts & techn. for hadron acc.

Coordination:
A. Lehrach/FZJ,
P. Spiller/GSI

ST3: ps – fs electron and photon beams

Coordination:
H. Schlarb/DESY,
A.-S. Müller/KIT


ST4: Novel acceleration concepts

Coordination:
U. Schramm/HZDR,
F. Grüner/U-Hamburg

Networking, workshops, joint projects and usage of infrastructure, transfer of new technologies between centers

Cooperation with German universities, international partners and industry

Cost per sub-topics



Coordination: Reinhard Brinkmann
Deputy: Andreas Jankowiak



SINBAD

horizontale, negative, ultraschnelle & accelerators at DESY

Kompakte Atto-Sekunden Lichtquelle
50 as, ICS
ERC Synergy
Grant, DESY,
Uni HH, Arizona

Ultraschnelle Elektronenpulse
< 1 fs mit konventioneller
Technologie
ARD, DESY, Uni
HH, KIT

Nutzbereichstudien Plasma-
beschleuniger, Skalierbarkeit
> 1 GeV/m, nutzbare Strahlqualität, FEL?
LAOLA, ARD, DESY, Uni HH

Raum für weitere Phasen
und Nutzer
Drittmitter
Interessengruppen
ELI

durch optimale Nutzung der Infrastruktur

ARD Spitzenforschung im alten DORIS Komplex

PIER

Ralph Assmann | SINBAD | 24.01.2014 | Page 16

R. Assmann, DESY

Jülich Short-Pulse Particle and Radiation Centre

Particle physics

Synchrotron radiation

JuSPARC

Material research

Markus Büscher

M. Büscher, FZJ

The LIGHT test-stand at GSI: coupling of laser-accelerated ions into conventional accelerators

Principle: manipulation of laser-accelerated ions

1. Laser-driven ion acceleration
2. beam conditioning (collimation)
3. drift line and phase-space rotation

Current results:

- Initial experimental proof of principle done
- Ion duration (high peak current) at 10 MeV energy
- diagnostics done
- in POF III
- Ion beam towards 100 MeV
- ions
- to GSI's SIS accelerator
- elements (repetition rate and

proton spectrum

proton beam

V. Bagnoud, GSI



bERLinPro centre for high power cw beams in sc accelerators

bERLinPro = Berlin Energy Recovery Linac Project
100 mA / low emittance technology demonstrator

beam dump
6.5 MeV, 100 mA = 650 kW

linac module
44 MeV

merger dogleg

booster
4.5 MeV

rf-gun
1.5-2 MeV

high virtual beam power zone
(microwave instability driven radiation generation)

50MeV, 100mA, 2ps (5 MW of virtual beam power)
50MeV, 10mA, <100fs (500kW of virtual beam power)
both modes normalized emittance < 1mm mrad

A. Jankowiak, HZB

ELBE center for high power radiation sources

Dual beam Petawatt / 150 TW ultrashort pulse laser facility

Diode pumped Petawatt laser development

Synchronized operation with ELBE accelerator

Dedicated shielded target areas (~1000m² laser lab space)

Beam driven sources (THz, FEL, ...) at ELBE

ELBE

150TW

Draco PW

HZDR

U. Schramm, HZDR

FLUTE: ARD-Forschung am KIT

- Ultrakurze Elektronenpulse (1 fs bis 300 fs)
- Grosser Bereich an Ladungen (1 pC bis 3 nC)
- Kohärente Strahlung für Materialwissenschaften und biologische Anwendungen
- Entwicklung/Tests von Kurzpuls-Strahlendiagnose und Instrumentierung
- Kooperation KIT, PSI, DESY

Ferninfrarot Linac- und

FLUTE

A. S. Müller, KIT

Helmholtz-Institute Jena


Development and application of novel plasma diagnostics:

- few-fs and 1-µm resolution,
- first direct visualization of the laser-driven plasma wave in a laser-electron accelerator.

M. Schwab et al., Applied Phys. Lett (2013)
A. Sävert et al., submitted (2013)

M. Kaluza, HIJ

Proposal: Helmholtz Distributed ARD Test Facility – Germany



The preparation team:

*R. Assmann (DESY), V. Bagnoud (GSI),
M. Büscher (HZJ), A. Jankowiak (HZB),
M. Kaluza (HIJ), A.-S. Müller (KIT),
U. Schramm (HZDR)*

> Theme 1: **Collaboration**

Networking of existing research infrastructure.

> Theme 2: **Synergy**

Extension of facilities for common usage.

> Thema 3: **Leadership**

2 flagship projects for internationally leading research with the **aim of ultra-compact accelerators and radiation sources** (plasma acceleration major player).

Conclusions – General

- > Conventional **accelerators are masterpieces of technology**, operating close to physical hard limits.
- > New projects (plan A) **scale/extend existing technology** → excellent performance but **larger size and cost**.
- > Many of us believe that a **plan B HEP project for the 2030's** is a good idea. Should be smaller and lower cost.
- > Plasma accelerators **reduce acceleration lengths by factor 100 – 1000**. Excellent candidate plan B technology.
- > Plasma accelerators: **potential for novel ultra-compact FEL's**, radiation sources, medical imaging (ultra-fast).
- > Plasma accelerators not fundamentally limited. **Problems are difficult but of technical nature**.



Our mission: The compact accelerator for high power physicists!



... for high power physicists!



The size of the accelerator structure is correct, but we want to place a plasma accelerator structure there with 10 GeV acceleration!

Conclusions – Europe

- > Europe: **Increased investments in novel accelerator R&D**. Accelerator R&D in Germany recognized as **independent research area**. Conventional and novel acc. R&D together. HEP and photon science acc. R&D together.
- > Europe: **≈15 significant projects** in plasma acceleration. EuroNNAc: exchange info, develop common plans.
- > Right time is now to spend time and efforts on **developing plasma accelerator technology to user readiness**.
- > Efforts on **grouping our European efforts**: EuPRAXIA proposal, CILEX, Helmholtz Distributed ARD Test Facility, ...
- > We hope that our efforts will be flanked by **strengthened US efforts on accelerator R&D**. It will pay off for US science and for all of us.



Thank you for your attention...



CILEX Project, Plateau de Saclay, France

- The big French project.
- Laser: APOLLON, 150J / 15fs, 1 shot/min, 5-10 PW , 10^{23} W/cm² plus satellite lasers.
- 450 m² experimental area for plasma acceleration of e⁻. Possibility to extend into 200m long acc. tunnel.



A. Specka



F. Amiranoff





X-5 Project at LOA, France

- > Salle Jaune Laser: 70 TW, repetition rate 10 Hz, pulse duration of 30 fs
- > Goals:
 - Exploration of new laser plasma accelerator blueprints for the production of electron beams and very strong peak point currents.
 - These electron beams will be used in a variety of domains, including life science and materials science.
 - They will also be used to study the blueprint of compact free electron lasers (FEL) for the production of intense X and XUV beams.



European
Research
Council



Victor Malka



Researcher at CNRS and Lecturer

An excellence grant for LOA.




Victor Malka, a CNRS researcher and lecturer in the physics department at X, works at **ENSTA**, in a team that he set up in 2001 to study **laser-plasma particle acceleration**. In July 2008, he was awarded a **grant** by the European Research Council of **2.2 million euros**. The grant was awarded in two categories: junior and senior. It was in the second category that he was rewarded for his many scientific works and for his ability to create **new fields of research**.



COXINEL Project at SOLEIL, France

- > COXINEL: COherent Xray source INferred from Electrons accelerated by Laser.
- > Leader: Marie-Emmanuelle Couprie, SOLEIL
- > Goals:
 - COXINEL aims to demonstrate that, by using laser acceleration, it is possible to obtain the free-electron laser (FEL) amplification needed to develop more compact light sources.
 - FEL are the first tunable X-ray lasers and the most intense light sources in the X-ray energy domain.
 - COXINEL benefits from a very favorable environment at SOLEIL and more widely on the Paris-Saclay campus, particularly in terms of engineering, to turn the ideas and theories that physicists have devised into reality.
- > Closely connected to project X-5 in LOA.





G. Mourou

Coherent Amplification Network

Figure 1 | Principle of a coherent amplifier network. An initial pulse from a seed laser (1) is stretched (2), and split into many fibre channels (3). Each channel is amplified in several stages, with the final stages producing pulses of ~1 mJ at a high repetition rate (4). All the channels are combined coherently, compressed (5) and focused (6) to produce a pulse with an energy of >10 J at a repetition rate of ~10 kHz (7).

The future is fibre accelerators

Gerard Mourou, Bill Brocklesby, Toshiki Tajima and Jens Limpert

Could massive arrays of thousands of fibre lasers be the driving force behind next-generation particle accelerators? The International Coherent Amplification Network project believes so and is currently performing a feasibility study.





SPARC LAB, Frascati, Italy

- > Electron beam: 150 MeV, multi-bunch, bunch length below 300 fs, 200 pC, 1 μm norm. emittance
- > FLAME laser: Ti:Sa, chirped pulse amplification (CPA), 200 TW, 25 fs long, 10 Hz repetition rate, wavelength 800 nm
- > SPARC LAB is a multi-purpose user facility. Includes experiments on laser- and beam-driven plasma wakefield acceleration experiments. See example of laser-generated electrons.
- > Comb project is unique: resonant beam driven plasma wakefields.



M. Ferrario



LUND Laser Center, Sweden

- > Well established lab, since 1992.
- > Laser: 60 TW + 5 TW + 3 TW (three separate compressors). Emphasis on:
 - Stability (pulse-to-pulse)
 - Reliability (day-to-day)
 - Contrast (ex. plasma mirrors)
 - Beam pointing (active control)
 - Wavefront (adaptive optics)
- > Limited by manpower.




C.G. Wahlstroem

X-rays



Protons



Electrons



ELI Beamlines, Czech republic

- Laser: 10-15 fs duration, up to 10 PW. End stage: a few kJ in 15 fs (~200 PW) with low repetition rate (minute based).
- Might be the big player in some years.
- New techniques for medical image-display and diagnostics, radiotherapy, tools for new materials developing and testing, latest in X-ray optics, etc.
- Laser-accelerated, versatile electron and proton/ion source emitting in an unprecedented energy range (1 GeV to 100 GeV). Provide a major contribution for the development of future high-quality and low-cost proton sources for cancer therapy.



G. Korn



AWAKE experiment, CERN


- International collaboration with approved experiment at CERN beam.
- Driver: 450 GeV proton bunch, $1e11$, $3.5\mu\text{m}$ emittance, bunch length \gg plasma wavelength
- Modulation experiment:
 - The first parts of the proton bunch induce a plasma wakefield that starts self-modulating the later parts of the bunches.
 - Microbunching starts happening:
 - 1) Protons at focusing regions survive.
 - 2) Protons at defocusing regions get lost.
 - Surviving microbunches induce wakefields.
 - Accelerate injected electrons from several 10 MeV to GeV.



A. Caldwell




M. Wing



CALA Project, Munich, Germany

- Builds on expertise at MPQ and LMU.
- Successful in laser-driven generation of electron beams. See example with MPQ lasers.
- Ongoing: 63 M€ investment. Financed by State Bavaria and Germany.
- CALA completion: 2015...

S. Karsch





- SCAPA = Scottish Centre for the Application of Plasma-based Accelerators
- LWFA for generation of particle beams (electrons, protons, neutrons and light ions) and radiation pulses (THz, infrared, X-rays and gamma rays).
- Experimental and theoretical research into high-power laser-plasma interactions.
- ALPHA-X Project: A Laser Wakefield Accelerator Dedicated to the Production and Application of Ultra-short Electron Bunches and Radiation Pulses.



D. Jaroszynski




B. Hidding



Z.M. Sheng

STFC – CLF and John Adams Institute in UK

- STFC Central Laser Facility used also for LWFA.
- John Adams Institute: Imperial College London, Oxford University, Royal Holloway, University of London.
- Plasma accelerators and novel sources of coherent x-rays. LWFA, narrow energy spread. Resonant LWFA. Medical imaging.
- Training of accelerator specialists, also in advanced methods.



S. Hooker



Z. Najmudin



A. Seryi



R. Walczak

Cockcroft Institute and CLARA Project

- Cockcroft Institute: Universities of Lancaster, Liverpool and Manchester, the Science and Technology Facilities Council (STFC at the Daresbury and Rutherford Appleton Laboratories), North West Development Agency (NWDA).
- Ultra-short pulse generation. FEL design and industrial applications → CLARA project. Plasma acceleration.
- Training.



J. Clarke



C. Welsch



S. Smith



> Laser: Ti:Sa 200 TW, 25 fs pulse length, 5 Hz repetition rate

- Initially: Laser-driven wakefields in REGAE. LUX exp. towards FEL.
- Later: Move to SINBAD facility.

> Beams:

- **REGAE**: 5 MeV, fC, 7 fs bunch length, 50 Hz



F. Grüner

A. Maier

- **FLASH**: 1.25 GeV, 20 – 500 pC, 20 - 200 fs bunch length, 10 Hz.

Beam-driven plasma wakefields. Beam-driven plasma wakefields with shaped beams and innovative injection methods. Helmholtz VI with UK collaboration.

FLASHForward ▶▶

- **PITZ**: 25 MeV, 100 pC, 20 ps bunch length, 10 Hz. Beam modulation experiment in a plasma cell, preparation to CERN experiment AWAKE



F. Stephan



J. Osterhoff

- Future **SINBAD**: 100 MeV, 0.01 – 3 pC, down to 10 fs bunch length, 10 – 1000 Hz

ARD top class research in existing infrastructure (discontinued DORIS)

Compact Atto-Second Light Source

50 as, ICS
 ERC Synergy Grant 14 M€, DESY, Uni HH, Arizona




European Research Council
 Established by the European Commission

P. Fromme F. Kärtner



R. Assmann H. Chapman



Ultrashort electron pulse
 < 1 fs with conventional technology
 ARD, DESY, Uni HH, KIT

Develop ability to use plasma acceleration, scalability
 > 1 GeV/m, useable beam quality, FEL?
 LAOLA, ARD, DESY, Uni HH

Footprint: 90 m x 50 m

Room for further phases and users
 Third party funding, interest from ELI, ...

